

Incidental Liver Findings in ^{99m}Tc -Sestamibi Myocardial Perfusion SPECT

Andrea De Lorenzo, Pedro Gomes da Cruz Netto, and Paulo de Sa Villela Pedras

Universidade Federal do Rio de Janeiro, Rio de Janeiro, RJ Brazil

In cardiac nuclear imaging, the liver is mostly remembered as a source of artifact in the left ventricular inferior wall. However, the liver may eventually display unexpected findings, which draw attention to significant noncardiac disease. In a patient referred for myocardial perfusion SPECT to investigate atypical chest pain, focal areas of increased tracer uptake revealed intrahepatic biliary dilatation.

Key Words: myocardial perfusion SPECT; ^{99m}Tc sestamibi; hepatobiliary; diagnosis

J Nucl Med Technol 2016; 44:42–43

DOI: 10.2967/jnmt.115.158451

Although the use of nuclear techniques for the diagnosis of biliary disorders is not new, as cholescintigraphy was once considered the noninvasive procedure of choice for the evaluation of biliary tract function, nuclear imaging of the biliary system is nowadays left for selected cases (1). The biodistribution of ^{99m}Tc -sestamibi, with accumulation in the liver (2), makes the hepatic image in myocardial perfusion SPECT a common source of artifact, creating either hot spots or false perfusion defects in the left ventricular inferior wall (3). On the other hand, this biodistribution enables the detection of incidental hepatobiliary abnormalities.

CASE REPORT

A 63-y-old woman with atypical chest pain was referred for myocardial perfusion SPECT. She reported prior surgical intervention in the pancreas due to pancreatitis secondary to severe hypertriglyceridemia. No follow-up regarding the abdominal intervention had been performed for several years.

A same-day, rest–dipyridamole stress (370 MBq/1,110 MBq) ^{99m}Tc -sestamibi myocardial perfusion SPECT study was performed. Images were obtained in a dual-head camera (Millenium MG; GE Healthcare). Rest acquisition

started 30 min after tracer injection. Postdipyridamole images were obtained 45 min after injection. Figure 1 shows scintigraphic findings. There were several areas of increased focal uptake in the liver, both at rest and after stress. Myocardial perfusion was considered overall normal, with a mild anteroapical perfusion defect, which was interpreted as breast attenuation in view of normal thickening and wall motion.

After the nuclear scan, further investigation of the liver abnormalities was indicated. The patient underwent abdominal CT scanning (Fig. 2), which revealed multiple intrahepatic biliary tract dilatations. The focal areas of increased tracer uptake in the scintigraphic images were therefore attributed to bile stasis within biliary tract dilatations. The patient was therefore referred for specialized, surgical follow-up. Laboratory evaluation showed serum bilirubin values within the reference range (total, direct, and indirect bilirubin of 1.2, 0.1, and 1.1, respectively); amylase, 21 U/L; lipase, 36 U/L; alkaline phosphatase, 779 U/L; γ glutamyl transferase, 659 U/L; fasting glucose, 359 mg/dL; and triglycerides, 2,740 md/dL. Because the patient was stable in relation to the hepatobiliary condition, no further intervention was performed. Her follow-up was noteworthy only for frequent episodes of severe hyperglycemia needing admission for glycemic control despite the domiciliary use of insulin.

DISCUSSION

The case illustrates the importance of careful inspection of raw images when analyzing myocardial perfusion




FIGURE 1. ^{99m}Tc -sestamibi myocardial perfusion SPECT images showing several areas of increased uptake in liver. (Left) Poststress images. (Right) Rest images.

Received Mar. 28, 2015; revision accepted May 1, 2015.
For correspondence or reprints contact: Andrea De Lorenzo, Av Brigadeiro Trompowsky s/n- Ilha do Fundão, Rio de Janeiro, RJ Brazil, 21941-913.
E-mail: andlorenzo@hotmail.com
Published online Jun. 25, 2015.
COPYRIGHT © 2016 by the Society of Nuclear Medicine and Molecular Imaging, Inc.




FIGURE 2. Abdominal CT scan showing multiple dilations of intrahepatic biliary tract (arrows). Liver had normal size and parenchyma, whereas gallbladder was not visualized. Biliodigestive bypass was found, with anastomosis of common bile duct to duodenum (not shown in figure).

scintigraphy. Eventually, extracardiac findings may point toward significant subclinical disease. In this case, the biodistribution of ^{99m}Tc -sestamibi enabled the detection of

biliary abnormalities, because tracer excretion into the biliary system occurs as long as liver uptake of ^{99m}Tc -sestamibi decreases, 30–60 min after injection (2,4).

CONCLUSION

In cardiac nuclear imaging, attention must be given to extracardiac findings, as these may uncover relevant pathology. Early diagnosis of disease may have important clinical implications, with impact on the patient's prognosis.

DISCLOSURE

No potential conflict of interest relevant to this article was reported.

REFERENCES

1. Weissmann HS, Frank MS, Rosenblatt R, Sugarman LA, Freeman LM. Role of ^{99m}Tc -IDA cholescintigraphy in evaluating biliary tract disorders. *Gastrointest Radiol.* 1980;5:215–223.
2. Wackers FJ, Berman DS, Maddahi J, et al. Technetium-99m hexakis 2-methoxyisobutyl isonitrile: human biodistribution, dosimetry, safety, and preliminary comparison to thallium-201 for myocardial perfusion imaging. *J Nucl Med.* 1989;30:301–311.
3. King MA, Xia W, de Vries DJ, et al. A Monte Carlo investigation of artifacts caused by liver uptake in single photon emission computed tomography perfusion imaging with technetium 99m labeled agents. *J Nucl Cardiol.* 1996;3: 18–29.
4. Iskandrian AS, Verani MS, Heo J. Pharmacologic stress testing: mechanism of action, hemodynamic responses, and results in detection of coronary artery disease. *J Nucl Cardiol.* 1994;1:94–111.